

Access this article on  
[ScienceDirect](#) ▶

## Alanine transaminase as a single marker is not reliable for selecting candidates of pegylated interferon and ribavirin treatment

[Heidar Sharafi](#) , [Seyed Moayed Alavian](#)

Middle East Liver Disease (MELD) Center, Tehran, Iran

[Maryam Keshvar](#) 

Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran

Received: April 3, 2014; Published Online: May 19, 2014

DOI: <http://dx.doi.org/10.1016/j.jfma.2014.04.002> |  CrossMark

Article Info

Abstract

Full Text



References

We read with great interest the article by Lin et al.<sup>1</sup> They studied the efficacy and safety of pegylated interferon  $\alpha$ -2a and ribavirin in 12 hemophilic patients with chronic hepatitis C. The current study is appreciable since hemophilic patients are considered as a special patients group and the data regarding the treatment of hepatitis C in these individuals are limited in the literature.

Conflicts of interest: The authors have no conflicts of interest relevant to this article.

© 2014 Published by Elsevier Inc.

### Article Tools

 [PDF \(185 KB\)](#) [Email Article](#) [Add to My Reading List](#) [Export Citation](#) [Create Citation Alert](#) [Cited by in Scopus \(0\)](#) [Request Permissions](#) [Order Reprints](#)  
(100 minimum order)